

Nonlinear Funct. Anal. Appl., 8, N2, (2003).

On the discrepancy principle

A.G. Ramm

Mathematics Department, Kansas State University,
Manhattan, KS 66506-2602, USA
ramm@math.ksu.edu

Abstract

A simple proof of the convergence of the variational regularization, with the regularization parameter, chosen by the discrepancy principle, is given for linear operators under suitable assumptions. It is shown that the discrepancy principle, in general, does not yield uniform with respect to the data convergence. An a priori choice of the regularization parameter is proposed and justified for some nonlinear operator equations.

1 Introduction.

In this note the convergence of the variational regularization with the regularization parameter chosen by the discrepancy principle is discussed for linear operators. In [1] and [2] one finds the references related to the discrepancy principle.

Our assumptions in this Section are:

i) A is a linear, bounded, injective operator on a Hilbert space H , the equation $A(y) = f$ has a solution y , A^{-1} is not continuous, so that (1.1) is an ill-posed problem, $\|f_\delta - f\| \leq \delta$, $\|f_\delta\| > \delta$.

Given $\{\delta, f_\delta\}$, one wants to approximate y stably in the sense

$$\lim_{\delta \rightarrow 0} \|R_\delta f_\delta - y\| = 0, \quad (1.1)$$

where R_δ is an operator defined on all of H , a regularizer. Let $F(u) := \|Au - f_\delta\|^2 + a\|u\|^2$, where $a > 0$ is a parameter, and let u_a be the solution to $F(u_a) = \inf F(u)$. Existence and uniqueness of this variational problem are known: if $B := A^*A$, then

$$u_a = (B + a)^{-1} A^* f_\delta, \quad (B + a)u_a = A^* f_\delta, \quad (1.2)$$

equation (1.2) is the Euler equation for F , it has a unique solution u_a since $B \geq 0$ and $a > 0$, and

$$F(u_a) \leq F(u), \quad \|Au_a - f_\delta\|^2 + a\|u_a\|^2 \leq \delta^2 + a\|y\|^2. \quad (1.3)$$

Assume $f_\delta \perp N(A^*)$, where $N(A^*)$ is the null-space of A^* . This is done without loss of generality, since f_δ enters under the sign of A^* in (1.2). The discrepancy principle (DP) (introduced by Morozov, see [1] and [2]) chooses $a = a(\delta)$ as the root of the equation:

$$h(a, \delta) := \|Au_a - f_\delta\| = C\delta, \quad C = \text{const} \geq 1, \quad \|f_\delta\| > \delta. \quad (1.4)$$

Note that $N(Q) = N(A^*)$, where $Q := AA^* \geq 0$, and

$$\|Au_a - f_\delta\| = \|[A(B + a)^{-1}A^* - I]f_\delta\| = \|Q(Q + a)^{-1} - I\|f_\delta\| = a\|(Q + a)^{-1}f_\delta\|,$$

where the known formula $(B + a)^{-1}A^* = A^*(Q + a)^{-1}$ was used.

Thus,

$$h = a\|(Q + a)^{-1}f_\delta\|.$$

This implies that, for a fixed pair $\{f_\delta, \delta\}$, the function $h(a, \delta)$ in (1.4) is a monotone increasing function of a on $(0, \infty)$, $\lim_{a \rightarrow 0} h(a, \delta) = 0$, where we have used the assumption $f_\delta \perp N(A^*)$, and $h(\infty, \delta) = \|f_\delta\| > \delta$, so that (1.4) has a unique solution $a = a(\delta)$, $\lim_{\delta \rightarrow 0} a(\delta) = 0$.

Define $R_\delta f_\delta := u_\delta := u_{a(\delta)}$, where $a(\delta)$ is given by the DP.

Theorem 1.1. *If A is compact, i) holds, and $R_\delta f_\delta := u_\delta$, then (1.1) holds.*

Proof. From (1.4) and (1.3), with $a = a(\delta)$ and $u_a := u_\delta$, one gets

$$\|u_\delta\| \leq \|y\|. \quad (1.5)$$

Thus, $u_\delta \rightharpoonup u$ (weak sequential convergence) as $\delta \rightarrow 0$, and $Au_\delta \rightarrow Au$, because A is compact. From (1.3), with $\delta \rightarrow 0$ and $a \rightarrow 0$, it follows that $Au = f$. By the injectivity of A , one gets $u = y$, so $u_\delta \rightharpoonup y$ as $\delta \rightarrow 0$. To prove that

$$\lim_{\delta \rightarrow 0} \|u_\delta - y\| = 0, \quad (1.6)$$

it is sufficient to prove that $\lim_{\delta \rightarrow 0} \|u_\delta\| = \|y\|$, and this follows from (1.5) because $\|y\| \leq \liminf_{\delta \rightarrow 0} \|u_\delta\| \leq \limsup_{\delta \rightarrow 0} \|u_\delta\| \leq \|y\|$. The first inequality is the lower semicontinuity of the norm in H , and the last one follows from (1.5). Theorem 1.1 is proved. $\square$

An alternative proof of (1.6) is based on the inequality

$$\|u_\delta - y\|^2 \leq 2\Re(y, y - u_\delta), \quad (1.7)$$

which follows from (1.5) easily. Since $u_\delta \rightharpoonup y$ as $\delta \rightarrow 0$, (1.7) implies (1.6). $\square$

2 Generalizations.

2.1. *One can drop the injectivity of A assumption and still get (1.6):*

If $N := N(A)$ is the null-space of A , $Ay = f$, and $y \perp N$, then $u_\delta := u_{a(\delta)}$, defined in (1.2), has the property $u_\delta \perp N$. Indeed, if $\phi \in N$, then $(u_\delta, \phi) = (f_\delta, A(B + a)^{-1}\phi) = (f_\delta, (Q + a)^{-1}A\phi) = 0$. Thus $\lim_{\delta \rightarrow 0} u_\delta = u \perp N$. Since the problem

$$Au = f, \quad u \perp N \quad (2.1)$$

has only one solution, it follows that $u = y$. We have proved:

Theorem 2.1. *If $N \neq \{0\}$ and y solves (2.1), then (1.6) holds.*

2.2. *Let us drop both the compactness of A and the injectivity of A assumptions.*

Theorem 2.2. *If A is a linear bounded operator, and y solves (2.1), then (1.6) holds.*

Proof. As above, one gets (1.5) with $u_\delta := u_{a(\delta)}$ defined by (1.2). Thus, $u_\delta \rightarrow u$ and $(B + a)u_\delta \rightarrow A^*f$ as $\delta \rightarrow 0$, and this implies $Bu = A^*f$. Indeed, $B \geq 0$ is a monotone continuous operator, and therefore it is weakly closed, i.e., $u_\delta \rightarrow u$ and $Bu_\delta \rightarrow g$ imply $Bu = g$. Recall that $a = a(\delta) \rightarrow 0$ and u_δ is bounded as $\delta \rightarrow 0$, so $\lim_{\delta \rightarrow 0} a(\delta)u_\delta = 0$. Since $f = Ay$, one has $B(u - y) = 0$, and, since $N(B) = N(A)$, one gets $Au = Ay = f$. We have already proved that $u \perp N$, thus $u = y$, and $u_\delta \rightarrow y$ as $\delta \rightarrow 0$. This and (1.7) imply (1.6). $\square$

2.3. *Let us drop the linearity of A assumption and propose an a priori choice of the regularization parameter.*

Consider the functional (cf [3]):

$$F(u) := \|Au - f_\delta\| + \delta\|u\|_1, \quad D(F) = H_1, \quad (2.2)$$

where $D(F)$ is the domain of F , the norm $\|\cdot\|_1$ is a norm of a Hilbert space H_1 , which is dense in H , complete, and the imbedding $i : H_1 \rightarrow H$ is compact. Let

$$m = m(\delta) := \inf_u F(u),$$

and let u_δ be any element such that

$$F(u_\delta) \leq m + \delta. \quad (2.3)$$

Theorem 2.3. *Assume that A is a nonlinear, continuous, injective map, A^{-1} is not continuous, $f = A(y)$, $y \in H_1$, and (2.3) holds. Then (1.6) holds.*

Proof. Let $F(u_n) \rightarrow m$. Then for all sufficiently large n one has:

$$m \leq \|A(u_n) - f_\delta\| + \delta\|u_n\|_1 \leq \delta + \delta\|y\|_1 := c_1\delta. \quad (2.4)$$

Choose $u_\delta := u_n$ such that (2.3) holds, and let $\delta \rightarrow 0$. Since $\|u_\delta\|_1 \leq c_1$, one has $u_\delta \rightarrow u$, $A(u_\delta) \rightarrow A(u)$, and (2.4) shows that $A(u) = f$. By the injectivity of A , one gets $u = y$, so (1.6) holds. $\square$.

3 DP does not yield uniform with respect to f convergence.

In this Section the notations of Section 1 are used.

Theorem 3.1 *Let $S_\delta := \{v : \|Av - f_\delta\| \leq \delta\}$, $u_\delta := u_{a(\delta)}$, $a(\delta)$ solves (1.4), A satisfies assumptions i) of Section 1, and $N(A^*) = \{0\}$.*

Then there are f_δ such that

$$\lim_{\delta \rightarrow 0} \sup_{v \in S_\delta} \|u_\delta - v\| \geq c = \text{const} > 0. \quad (3.1)$$

Proof. Denote $(B + a)^{-1}A^* := T$, $B := A^*A$, and $u_\delta = Tf_\delta$. Then $\|T\| = \frac{1}{2\sqrt{a}}$ (see, e.g., [1]). Thus, there exists a $p = p_a$, such that

$$\|Tp\| \geq \frac{\delta}{8\sqrt{a}}, \quad \|p\| = 0.5\delta.$$

Let $R(A)$ denote the range of A . Since $R(A)$ and $R(B)$ are dense in H , and $N(A^*) = \{0\}$, there is a $z = z_a$ such that

$$\|f_\delta - AB^bz - p\| \leq \frac{\delta}{8}, \quad b = \text{const} \in (0, 1) \quad \|p\| = 0.5\delta. \quad (3.2)$$

For any v , one has:

$$\|Tf_\delta - v\| \leq \|Tf_\delta - TA v\| + \|TA v - v\|. \quad (3.3)$$

If $v = B^bz$, then

$$\lim_{\delta \rightarrow 0} \sup_{v \in S_\delta, v=B^bz, \|z\| \leq M} \|TA v - v\| = 0, \quad (3.4)$$

where $M > 0$ is an arbitrary large constant, and we have used the following formulas:

$$TA v - v = [(B + a)^{-1}B - I]B^bz = -a(B + a)^{-1}B^bz,$$

$$\|a(B + a)^{-1}B^b\| = a \sup_{0 \leq s \leq \|B\|} \frac{s^b}{s + a} = ca^b, \quad c = b^b(1 - b)^{1-b}.$$

From (3.3) and (3.4) it follows that

$$\lim_{\delta \rightarrow 0} \sup_{v \in S_\delta, v=B^bz, \|z\| \leq M} \|Tf_\delta - v\| = 0 \quad (3.5)$$

if and only if

$$\lim_{\delta \rightarrow 0} \sup_{v \in S_\delta, v=B^bz, \|z\| \leq M} \|Tf_\delta - TA v\| = 0. \quad (3.6)$$

Choose $v = B^bz$, where $z := z_a$. Then, using the triangle inequality and (3.2), one gets:

$$\|f_\delta - Av\| \leq \|p\| + \frac{\delta}{8} \leq \delta, \quad (3.7)$$

and

$$\|Tf_\delta - TA v\| = \|T(f_\delta - Av - p) + Tp\| \geq \|Tp\| - \frac{\delta}{8}\|T\| \geq \frac{\delta}{\sqrt{a}}\left(\frac{1}{8} - \frac{1}{16}\right) = \frac{\delta}{16\sqrt{a}}. \quad (3.8)$$

If one finds f_δ for which equation (1.4) has a root

$$a = c_1 \delta^2 [1 + o(1)], \quad c_1 = \text{const} > 0, \quad \delta \rightarrow 0, \quad (3.9)$$

then $\frac{\delta}{\sqrt{a}} \geq c = \text{const} > 0$, and (3.8) implies that (3.6) fails, so (3.5) fails, and Theorem 3.1 follows.

There are infinitely many such f_δ . Let us construct one of them. Assume, for simplicity, that $A = A^*$, so $B = A^2$, and (1.4) is:

$$a^2 \sum_{j=1}^{\infty} \frac{g_j}{(\mu_j + a)^2} = C^2 \delta^2, \quad g_j := |f_{\delta j}|^2, \quad \sum_{j=1}^{\infty} g_j < \infty, \quad (3.10)$$

where $\mu_j > 0$ are the eigenvalues of A^2 . Let $g_j = j^{-2}$, $\mu_j = j^{-1}$. Then, as we prove below,

$$\phi := \sum_{j=1}^{\infty} \frac{j^{-2}}{(j^{-1} + a)^2} = a^{-1} [1 + O(a)] \quad a \rightarrow 0. \quad (3.11)$$

If (3.11) holds, then (3.10) yields (3.9), and the argument is completed.

To prove (3.11), define $\psi := \int_1^\infty x^{-2}(x^{-1} + a)^{-2} dx$, and, setting $t = x^{-1}$, one gets

$$\psi = \int_0^1 (t + a)^{-2} dt = a^{-1} [1 + O(a)] \quad a \rightarrow 0.$$

This and the estimate

$$\phi - (1 + a)^{-2} < \psi < \phi,$$

yield (3.11). $\square$

In our construction, $w := f_\delta \notin R(A)$ does not depend on δ , and $f = Av_\delta$ is chosen for each $\delta > 0$ in a δ -neighborhood of f_δ (see (3.7)), so that f depends on δ . The element $v_\delta = B^b z_a$, $a = a(\delta)$, cannot converge as $\delta \rightarrow 0$: if $v_\delta \rightarrow v$ and $Av_\delta \rightarrow w$ then, by the continuity of A , one has $Av = w$. This is a contradiction, since $w \notin R(A)$. This contradiction proves that v_δ cannot converge. If A is compact, then a similar argument proves that $\lim_{\delta \rightarrow 0} \|v_\delta\| = \infty$.

References

- [1] Engl H., Hanke M., Neubauer A., Regularization of Inverse Problems, Kluwer, Dordrecht, 1996.
- [2] Morozov V., Methods of Solving Incorrectly Posed Problems, Springer Verlag, New York, 1984.
- [3] Ramm, A.G., Regularization of ill-posed problems with unbounded operators, J. Math. Anal. Appl., 271, (2002), 547-550.